



Cimarron[®] PLUS

Season-long control,

for more successful results //

Cattle and forage producers have wanted and needed a herbicide to help manage problem weeds in rangeland and pastures. They are looking for solutions that are economical and have no grazing or haying restrictions. Cimarron[®] PLUS delivers on all three. It offers residual control to protect pasture and rangeland grasses from problem broadleaf weeds, and it offers good activity on woody species.

About Cimarron PLUS

Cimarron PLUS delivers broad-spectrum weed control with one application. It offers significant flexibility; it can be tank-mixed with other pasture herbicides and applied by ground or aerial equipment, or as a spot treatment. There is no waiting period for grazing. Plus, it works quickly. Effects may be seen within 1-3 weeks.

Application and use

- // Broad-spectrum herbicide for the control and suppression of broadleaf weeds and brush
- // For best results, apply as a broadcast spray to young, actively-growing weeds from spring to early summer (see label for rates)
- // A spray adjuvant must be used in the spray mix unless otherwise specified on the label
- // Cimarron PLUS is registered for use on land dedicated to the production of grass forage in:
 - Rangeland
 - Pastures
 - Grass hay fields
 - Grasses in the Conservation Reserve Program (CRP)
 - Newly-seeded pastures (pre-plant, pre-emerge & post-plant)
 - Perennial weeds and grasses

Application tips

- // Spring spray applications help provide residual control of late-germinating weeds
- // For ground-based broadcast applications: apply using flat-fan nozzles and a minimum of 10 gallons per acre carrier volume
- // For aerial broadcast applications: use a minimum of 3 gallons per acre carrier volume
- // For spot applications: add 2-4 pints of non-ionic surfactant per 100 gallons of water

CAUTION: For use on grass pastures and rangeland only. Legumes such as alfalfa and clover are highly sensitive to Cimarron PLUS and may become stunted and/or die. Refer to the product label for important crop rotation information. Always follow label directions before applying to fescue or timothy.

Benefits of using Cimarron PLUS:

- // No grazing restrictions
- // Controls musk thistle – even if it is bolting, eliminating this noxious weed
- // Systemic action – translocates through the plant, including the root system
- // Easy to store, handle, measure and mix
- // May be tank-mixed with other pasture herbicides
- // Legume recrop flexibility
- // Not a federally restricted-use pesticide

Perennial weeds and woody plants controlled or suppressed:

- // Annual Broomweed
- // Thistle Complex
- // Bahiagrass
- // Marshelder
- // Sericea Lespedeza
- // Dewberries/Blackberries
- // Henbit
- // And more

Please see the product label for a full list of weeds and brush controlled. Do not use on crops.

For more information, contact your local Bayer representative or visit vm.bayer.us.

Do not use Cimarron PLUS in the following counties of Colorado: Alamosa, Conejos, Costilla, Rio Grande and Saquache.

ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS.

Bayer Environmental Science, a Division of Bayer CropScience LP, 5000 CentreGreen Way, Suite 400, Cary, NC 27513. For additional product information, call toll-free 1-800-331-2867. www.environmentalscience.bayer.us. Not all products are registered in all states. Bayer, the Bayer Cross and Cimarron are registered trademarks of Bayer. ©2019 Bayer CropScience LP. ES-1118-VM-CMRN-002-A-R0



Range &
Pasture