



Rezilon™ Herbicide

Break free

from ineffective grass control



New Rezilon™ (indaziflam) herbicide provides consistent control of summer and winter annual grasses and broadleaf weeds with residual activity and a wider application window, allowing you to increase hay quality and production.

Long-term control with Rezilon results in:

- // Higher quality hay in first harvest
- // Extended residual weed control
- // Increased fertilizer utilization because of fewer weeds present
- // Control of resistant ryegrass
- // Less drying time for cut hay

Application Chart

Weed Target	Timing	Rate
Crabgrass, Goosegrass, foxtails, and other warm-season annuals	January-March	3-5 fl oz/A
Ryegrass, little barley, and other cool-season annuals	August-September	3-5 fl oz/A
All susceptible weeds for year-long control	January-March August-September	3 fl oz/A 3 fl oz/A

When weeds have already germinated or perennial species are present, an appropriate tank-mix partner should be added for postemergence control. When tank-mixing, follow the correct mixing order of all products going into the spray tank. Dry products go in first, then emulsifiable concentrates, then Rezilon, followed by soluble products, then adjuvants. If tank-mixing Roundup® and Rezilon for a late-winter burndown and early preemergence treatment, add Rezilon first and allow enough time and agitation for thorough mixing, then Roundup, followed by the adjuvant. If correct mixing order is not observed, spray incompatibility and reduced weed control may result.

Rezilon benefits:

- // **Wider application window:**
Applications can be started as early as August and into the following spring as a result of extended residual weed control.
- // **Consistent weed control:**
Provides pre-emergence control of annual grass and broadleaf weeds in warm-season Bermudagrass and Bahiagrass hayfield sites.
- // **Increased hay quality:**
When applied according to label directions, growers could see higher-quality forage on their first cut as a result of controlling annual weed species present.
- // **Long residual activity:**
Once applied, Rezilon herbicide can sit on the soil surface for some time with no degradation from sunlight while waiting on rainfall for activation.

Key weeds controlled:

Rezilon controls more than 60 broadleaf and annual grass weeds, including:

- // Sandbur*
- // Crabgrass
- // Ryegrass
- // Goosegrass
- // Annual foxtails

*Split applications/full rate needed for optimum control.

IMPORTANT: This bulletin is not intended to provide adequate information for use of these products. Read the label before using these products. Observe all label directions and precautions while using these products.

For more information visit www.rezilon.com.



ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS.

©2020 Bayer Group. Always read and follow label instructions. Bayer, the Bayer Cross, Rezilon and Roundup are trademarks of Bayer. Not all products are registered in all states. For additional product information, call toll-free 1-800-331-2867. www.environmentalscience.bayer.us. Bayer Environmental Science, a Division of Bayer CropScience LP, 5000 CentreGreen Way, Suite 400, Cary, NC 27513.

Tank mixtures: The applicable labeling for each product must be in the possession of the user at the time of application. Follow applicable use instructions, including application rates, precautions and restrictions of each product used in the tank mixture. Not all tank mix product formulations have been tested for compatibility or performance other than specifically listed by brand name. Always predetermine the compatibility of tank mixtures by mixing small proportional quantities in advance.